

n-Categories Admissible by n-graph*

Władysław Marcinek* and Zbigniew Oziewicz†

Submitted January 3, 2001

Abstract

We consider examples of the n-categories having the same n-graph.

Contents

1	n-category as n-graph with a composition	1
2	Graph of graphs and category of categories	4
3	n-Categories admissible by n-graph	8

1 n-category as n-graph with a composition

We define an n-category as an n-graph equipped with the partial compositions of cells, or even with the graftings of cells in case a family of 0-cells is a strict

*Submitted to ‘Miscellanea Algebraica’, Waldemar Korczyński and Adam Obtułowicz, Editors, Akademia Świętokrzyska, Kielce, Poland.

Supported by el Consejo Nacional de Ciencia y Tecnología (CONACyT) de México, proyecto # 27670 E (1999-2000), and by UNAM, DGAPA, Programa de Apoyo a Proyectos de Investigación e Innovación Tecnológica, proyecto # IN 109599.

¹Institute of Theoretical Physics, University of Wrocław, Plac Maksa Borna 9, 50-204 Wrocław, Poland, wmar@ift.uni.wroc.pl

²Universidad Nacional Autónoma de México, Facultad de Estudios Superiores, Apartado Postal 25, 54700 Cuautitlán Izcalli, Estado de México, oziewicz@servidor.unam.mx; and Uniwersytet Wrocławski, Instytut Fizyki Teoretycznej, Pl 50204 Wrocław, Poland.

A member of Sistema Nacional de Investigadores, México, No de expediente 15337.

monoid [Oziewicz & Vázquez Coutiño 1999]. Therefore here n -graph G is a carrier of n -category (G, c_G) and there is a forgetful functor from n -category to his n -graph, $(G, c_G) \mapsto G$.

Definition 1.1 (Graph). An ∞ -graph G is an infinite sequence of families of i -cells $\{G_i, i \in \mathbb{Z}\}$ a sequence of sourcees $\{s_G \equiv s_i\}$ and targets $\{t_G \equiv t_i\}$ surjections, and the family of sections $\{i_G \equiv i_i\}$,

$$\cdots G_0 \begin{array}{c} \xrightarrow{s_0} \\ \xleftarrow{i_0} \\ \xrightarrow{t_0} \end{array} G_1 \begin{array}{c} \xrightarrow{s_1} \\ \xleftarrow{i_1} \\ \xrightarrow{t_1} \end{array} G_2 \begin{array}{c} \xrightarrow{s_2} \\ \xleftarrow{i_2} \\ \xrightarrow{t_2} \end{array} \cdots$$

Let type $x \equiv (sx, tx)$. By definition all 0-cells are of the same type, there are no more then two (-1) -cells and at least must be one (-1) -cell, $G_{-1} \equiv \{s_{-1}G_0, t_{-1}G_0\}$. A bimap $G_i : G_i \times G_i \rightarrow 2^{G_{i+1}}$ is defined as follows. If $x, y \in G_i \times G_i$, then

$$G_i(x, y) \equiv \{z \in G_{i+1}, \text{type } z = (x, y)\} \subset G_{i+1}.$$

The following conditions must hold.

Type. If type $x \neq \text{type } y$, then $G_i(x, y) = \emptyset$.

Section. $s_i \circ i_i = \text{id}_{G_i} = t_i \circ i_i$.

Definition 1.2 (n -graph). If $\forall i \in \mathbb{N}$, $G_{i+n} \simeq G_n$, then an ∞ -graph G is said to be n -graph. Hence $s_{i+n} = t_{i+n}$ and $i_{n+i} \circ s_{n+i} = \text{id}_{G_{n+i+1}}$.

In the terminology of Street [1998, p. 100] what we call n -graph G is said to be n -skeletal globular object, however we do not adopt this terminology in what follows. For example we do not wish to exclude from the further consideration more then two surjections $\{s, t, \dots\}$.

Definition 1.3 (Skeletal graph). If $\forall x, y$, such that type $x = \text{type } y$ and $\forall i \in \mathbb{N}$, $|G_i(x, y)| = 1$, then a graph G is said to be skeletal.

Definition 1.4 (n -category). An n -graph G with a operation c_G , is said to be an n -category,

$$c_G \equiv \{G_i(x, y) \times G_i(y, z) \xrightarrow{c_i xyz} G_i(x, z)\}.$$

In case $c_i xyz$ needs not to be global $\forall x, y, z$ then n -category is said to be partial.

Steiner [1998] introduced ‘genuine’ and partial n -category as a n -category generated by a ‘complex’. Therefore the Steiner’s complex is generating a unique n -graph admitting a unique n -category (either partial or global. In our definition an n -graph do not need admit a unique n -category (either global or partial).

Comment 1.5. An n -category G is said to be *strict* if c_G is associative, *i.e.* if a theory is equational. Otherwise an n -category is said to be *weak*. Gray [1976] investigated tensor product of 2-categories. Simpson [1998] studied strict n -category, 3-grupoids and homotopy types.

A weak 2-category is said to be a bicategory [Bénabou 1967]. A weak 3-category is said to be a tricategory [Gordon, Power & Street 1995; Baez & Dolan 1996; Leinster 1998]. Every tricategory is triequivalent to a category of bicategories equipped with the strong product of Gray [1976]. Baez and Dolan [1996], Baez [1997] and Leinster [1998], considered a weak n -category.

Batanin introduced a globular category [Batanin 1998, Street 1998].

Terminology 1.6. 0-cell is also called an object, 0-morphism, node; 1-cell - a morphism, 1-morphism, an arrow, a functor (Definition 3.2). A 2-cell is called a 2-morphism, morphism of morphism, natural transformation (Definition 3.3), etc.

Lemma 1.7. A skeletal n -graph admits exactly one n -category.

Definition 1.8 (Operad = Sketch with graftings). Bénabou noticed in 1967 that G_0 is a monoid iff there is a single (-1) -cell only. Then a n -category G is said to be a monoidal. If G_0 is a *free* monoid on k -generators, then a category G is said to be a k -sorted sketch. In this case a partial operation c_G from Definition 1.4 generalize to grafting $G_1 \times G_1 \rightarrow 2^{G_1}$ [Oziewicz & Vázquez 1999].

The above observation by Bénabou suggest to encode further category structure in a hidden negative $\mathbb{Z}$ -grade ‘tail’ of a graph G as follows.

Definition 1.9 (Structure tail). A finite sequence of negative $\mathbb{Z}$ -grade cells $\{G_k, -\infty < k \leq -1\}$ with $|G_k| \leq 2$, is said to be a structure tail of the category G .

Definition 1.10 (Opposite category). Let C be n -category. Then $\forall i \geq 1$, i -opposite n -category C_i^{opp} is defined by ‘reversing’ i -cells, $s_i^{opp} = t_i$, $t_i^{opp} = s_i$, and unchanged j -cells for $j \neq i$, $s_{j \neq i}^{opp} = s_j$, $t_{j \neq i}^{opp} = t_j$.

Example 1.11. A collection $G_i(x, y) \subset G_{i+1}$ is a 1-category,

$$(G_i(x, y))_0 \equiv G_i(x, y),$$

$$(G_i(x, y))_1 \equiv \{[G_i(x, y)](\alpha, \beta) \subset G_{i+2} : \quad \forall \alpha, \beta \in G_i(x, y)\}.$$

Example 1.12. Let $\mathbb{k}$ be a field. Let every 0-cell be a $\mathbb{k}$ -space and thus G_0 be unital $\mathbb{N}$ -algebra (unital rig) with $\oplus$ and $\otimes \equiv \otimes_{\mathbb{k}}$. Let $G_0(V, W)$ be a $\mathbb{k}$ -space $\text{lin}_{\mathbb{k}}(V, W)$ and $G_0(V, W)(f, g) \subset G_1(f, g) \subset G_2$ with $|G_1(f, g)| = 1$. Then G is a 2-category.

Example 1.13 (Wolff 1974). Enriched Wolff’s graph = 2-graph.

Example 1.14. n -Grupoid is an n -category with an inversion [Simpson 1998].

Definition 1.15 (Cocategory). An n -graph G with a binary cooperation $\triangle_G$ is said to be an n -cocategory,

$$\triangle_G \equiv \left\{ G_i(x, y) \xrightarrow{\triangle_{i xy}} G_i(x, \dots) \times G_i(\dots, y) \right\}.$$

2 Graph of graphs and category of categories

In this Section G is ∞ - or n -graph such that G_0 is a collection of k -graphs, so G is a ‘ n -graph of k -graphs’.

Definition 2.1 (Covariant vs contra). Let $E, F \in G_0$ be two k -graphs. An k -graph (absolutely covariant) morphism $f \in G_0(E, F) \subset G_1$ is a sequence of mappings $\{f_i\}$ intertwining s_E, t_E, i_E with s_F, t_F, i_F ,

$$\begin{array}{ccccccc} E_0 & \xleftarrow{s_0, t_0} & E_1 & \xleftarrow{s_1, t_1} & E_2 & \xleftarrow{s_2, t_2} & \dots, \\ f_0 \downarrow & & f_1 \downarrow & & f_2 \downarrow & & f \downarrow \\ F_0 & \xleftarrow{s_0, t_0} & F_1 & \xleftarrow{s_1, t_1} & F_2 & \xleftarrow{s_2, t_2} & \dots \end{array}$$

A 1-cell $f \in G_0(E, F_i^{opp})$ or $\in G_0(E_i^{opp}, F)$ is said to be i -contravariant.

Weakening: if the above intertwining (= a commutativity $f \circ s = s \circ f$) is up to 2-cell $\alpha : f \circ s \rightarrow s \circ f$, then f is said to be pseudo-morphism or α -morphism, or weakened morphism.

Definition 2.2 (Functor). Let (E, c_E) and (F, c_F) be k - or ∞ -categories. An k -graph morphism $f \in G_0(E, F)$ is said to be an k -category morphism $f \in G_0(c_E, c_F)$ if f intertwine c_E with c_F ,

$$\begin{array}{ccc} E_i(x, y) \times E_i(y, z) & \xrightarrow{c_E} & E_i(x, z) \\ f_{i+1} \times f_{i+1} \downarrow & & f_{i+1} \downarrow \\ F_i(f_i x, f_i y) \times F_i(f_i y, f_i z) & \xrightarrow{c_F} & F_i(f_i x, f_i z) \end{array}$$

Definition 2.3 (Natural transformation, Eilenberg & Mac Lane 1945).

Let $f, g \in G_0(c_E, c_F)$. A k -transformation $t \in G_1(f, g)$, is a collection of $(i+1)$ -cells $tx \in F_i(f_i x, g_i x)$, intertwining f with g ,


$$\begin{array}{ccc} E_i(x, y) & \xrightarrow{f_{i+1}} & F_i(f_i x, f_i y) \\ g_{i+1} \downarrow & & (t_y)_* \downarrow \\ F_i(g_i x, g_i y) & \xrightarrow{(t_x)^*} & F_i(f_i x, g_i y) \end{array}$$

For $\forall a \in E_i(x, y)$, $(g_{i+1}a) \circ (tx) = (ty) \circ (f_{i+1}a) \in F_{i+1}$.

In the above Definitions 2.1-2.3 we used the (labelled) commutative diagrams as invented by Witold Hurewicz in 1941. In the next Definition we prefer to use the ‘graphical calculus’, *i.e.* the relations (also known as ‘identities’), in an operad of graphs. This calculus is has been invented among other by Yetter [1990], by Joyal & Street [1993], and can be also coined as the Yetter-Joyal-Street *string* calculus. Calculus of n -graphs is an alternative language for commutative diagrams. Hurewicz associated to i -cell from G_i the ‘ i -dimensional geometrical arrow’, an oriented solid, *i.e.* 0-dimensional (labelled) vertex, a point, as a 0-cell $\in G_0$, a 1-dimensional directed arrow as a 1-cell from G_1 , a 2-dimensional oriented surface as a 2-cell from G_2 , etc. The graphical calculus for an n -graph G consists in representing n -cell as a 0-dimensional vertex, then $(n-1)$ -cell is represented as a 1-dimensional directed line, an arrow, etc, and finally 0-cell from G_0 is represented as an oriented n -dimensional volume.


Any segment of two consequent families of n-graph G , $G_i \leftarrow G_{i+1}$, can be represented in the operad of graphs as follows: if one-dimensional directed edge (an arrow) represents an i -cell, then zero-dimensional node must represents an $(i+1)$ -cell, *i.e.* an $(i+1)$ -morphism of i -cells.



Any segment of three families of n-graph G can be represented in these alternative languages as follows



All graphs in an operad are implicitly directed from the top (top is an input) to the bottom (output). Dashed boxes (from infinity) are for convenience. For two families segments all nodes are inner, there are no outer nodes. For three families segments all nodes and arrows are inner, there are neither outer arrows, nor outer nodes [Oziewicz & Vázquez Coutiño 1999].

Because in our convention all graphs are implicitly directed from the top to the bottom, therefore what is known in the commutative diagrams of opetopes [Baez and Dolan 1997], globes [Batanin 1998], and so on, as the vertical composition, in an operad of graph is the horizontal composition rather and vice versa. The following Figures illustrate this correspondence.






Definition 2.4 (Modification, Bénabou 1967). Let $s, t \in G_1(f, g)$. A modification $\mu \in G_2(s, t)$ is a collection $(i+2)$ -cells $\mu x \in F_{i+1}(sx, tx)$, intertwining s with t , such that $\forall a, b \in E_i(x, y) \subset E_{i+1}$ and

$$\forall \alpha \in E_i(x, y)(a, b) \subset E_{i+1}(a, b) \subset E_{i+2},$$

the following equations hold $\forall i \in \mathbb{N}$,



$$\begin{aligned}
x, y \in E_i, \quad a, b \in E_{i+1}, \quad \alpha \in E_{i+2}, \\
\text{surfaces:} \quad fx, gx, fy, gy &\in F_i, \\
\text{arrows:} \quad sx, tx, sy, ty, fa, fb, ga, gb &\in F_{i+1}, \\
\text{nodes:} \quad \mu x, \mu y, f\alpha, g\alpha &\in F_{i+2}.
\end{aligned}$$

Each of the above graphs, two elements of an operad, contains four paths from fx to gy , bypassing over or under nodes. The above relation in an operad tells that these four paths must be equal,

$$\begin{aligned}
(ga) \circ (sx) &= (sy) \circ (fa), \\
(gb) \circ (sx) &= (sy) \circ (fb), \\
(ga) \circ (tx) &= (ty) \circ (fa), \\
(gb) \circ (tx) &= (ty) \circ (fb), \\
\text{and } "(g\alpha) \circ (\mu x) &= (\mu y) \circ (f\alpha)".
\end{aligned}$$

Therefore Definition 2.4 include Definition 2.3. Definition 2.4 is for absolutely covariant f and g .

Let $\mu, \nu \in G_0(c_E, c_F)(f, g)(s, t) \subset G_1(f, g)(s, t) \subset G_2(s, t) \subset G_3$.

A ‘modification of modifications’ $A \in G_3(\mu, \nu) \subset G_4$ is a collection of $(i + 3)$ -cells $Ax \in F_{i+2}(\mu x, \nu x)$, intertwining μ with ν such that the set of relations hold $\forall i \in \mathbb{N}$. These relations needs graphics with 3-dimensional volumes, this time $x, y \in E_i$ need to be represented by 3-dimensional volumes and Ax and Ay by nodes.

3 n-Categories admissible by n-graph

Problem 3.1. Given n-graph G . How many (non isomorphic) n-category structures are allowed?

Example 3.2. Let G be a 2-graph whose 0-cells are sets, 1-cells are mappings between sets, 2-cells are changes of mappings with the same domain and codomain. Then every 3-cell is trivial.

Example 3.3. A 2-graph of 1-categories is a 2-category whose 1-cells are functors and 2-cell are natural transformations.

Example 3.4. Let G be a 1-graph such that every 0-cell $x \in G_0$ is an $(n-1)$ -category, and whose i -cells are $(n-1)$ -category morphisms. If there is an (associative) composition

$$c = \{G_0(x, y) \times G_0(y, z) \xrightarrow{c_{x,y,z}} G_0(x, z)\}, \quad (1)$$

then we obtain an n -category G' with only one 0-cell G and whose i -cells are $(i-1)$ -cell of G .

Definition 3.5 (Cobordism of manifolds, Stong 1998). A manifold M is said to be a cobordism from A to B if exists a diffeomorphism from a disjoint sum, $\varphi \in \text{diff}(A^* \cup B, \partial M)$. Two cobordisms $M(\varphi)$ and $M'(\varphi')$ are equivalent if there is a $\Phi \in \text{diff}(M, M')$ such that $\varphi' = \Phi \circ \varphi$. The equivalence class of cobordisms is denoted by $M(A, B) \in \text{Cob}(A, B)$.

Example 3.6. The unit cobordism $\partial(\Sigma \times [0, 1]) = \Sigma \cup \Sigma^*$.

Nontrivial examples of cobordism are given by Ionićiu [1997].

The 1-graph of oriented cobordisms Cob has oriented manifolds of the fixed dimension as objects, Cob_0 , and Cob_1 are cobordisms. A Cob_0 is a Grothendieck rig $\equiv$ N-algebra, *i.e.* an commutative monoid with duality, a monoidal structure is given by direct union ‘ $\cup$ ’ and the duality is reversal of orientation.

Composition of cobordisms c_{Cob} comes from gluing of manifolds. Let $\varphi' \in \text{diff}(C^* \cup D, \partial N)$. One can glue cobordism M with N by identifying B with C^* , $(\varphi')^{-1} \circ \varphi \in \text{diff}(B, C^*)$. We obtain the glued cobordism $(M \circ N)(A, D)$ and a semigroup operation,

$$c(A, B, D) : \text{Cob}(A, B) \times \text{Cob}(B, D) \longrightarrow \text{Cob}(A, D).$$

Example 3.7. The 1-graph of oriented cobordisms Cob with the composition c_{Cob} is a 1-category.

A *surgery* is an operation of cutting a manifold M and gluing to cylinders. A surgery gives new cobordism: from $M(A, B)$ into $N(A, B)$. The disjoint sum of $M(A, B)$ with $N(C, D)$ is a cobordism $(M \cup N)(A \cup C, B \cup D)$. We got a 2-graph of cobordism Cob with $\text{Cob}_0 = \text{Man}_d$, $\text{Cob}_1 = \text{Man}_{d+1}$, whose 2-cells from Cob_2 are surgery operations.

Kerler [1998] found examples of categories formed by classes of cobordism manifolds preserving disjoint sum or surgery. These examples are discussed by Baez and Dolan [1996].

Example 3.8. Let G be an 2-graph such that every 0-cell is a k -graph, every 1-cell $f \in G_0(x, y)$ is a k -graph morphism, composition of 1-cells is defined as a composition of k -graph morphisms. If every 2-cell $t \in G_1(f, g)$ is a k -graph transformation intertwining f with g , where $f, g \in G_0(x, y)$, and every k -cell $k = 3, \dots$, is trivial, then the 2-graph G becomes a 2-category.

Example 3.9. Let G be an 3-graph such that every $x \in G_0$ be a k -category with a composition c_x . Let $f, g \in G_0(c_x, c_y)$ be a k -category morphisms, all 2-cells $s, t \in G_1(f, g)$ be a k -category transformations, and let every 3-cell $\mu \in G_2(s, t)$ be a modification. Compositions of k -category morphisms, transformations and modifications define compositions making the 3-graph G a 3-category.

References

- [1] Baez John C. and James Dolan (1996), Higher-dimensional algebra and topological quantum field theory, *Journal of Mathematical Physics* **36**, 6073
- [2] Baez John C. (1997), An Introduction to n-Categories, q-alg/9705009.
- [3] Batanin Michael A. (1998), Monoidal Globular Categories As a Natural Environment for the Theory of Weak n-Categories, *Advances in Mathematics* **136** 39–103
- [4] Bénabou Jean (1967), Introduction to bicategories, in: *Lecture Notes in Mathematics* **47** 1–17
- [5] Crane Louis & David N. Yetter (1994), On algebraic structures implicit in topological quantum field theories, *Journal of Knot Theory and Reimifications* (1995), hep-th/9412025
- [6] Crane Louis & David N. Yetter (1996), Examples of categorification, q-alg/9607028
- [7] Ehresmann C., *Catégories et Structures*, Dunod, Paris 1965
- [8] Eilenberg Samuel and Saunders Mac Lane, General theory of natural equivalence, *Transactions of American Mathematical Society* **58** (1945) 231–294

- [9] Freyd P. and David N. Yetter, Braided compact monoidal categories with applications to low dimensional topology, *Advances in Mathematics* **77** (1989) 156–182
- [10] Gray J. W. (1976), Coherence for the tensor product of 2-categories, in: *Algebra, Topology and Category Theory* (a collection of papers in honor of Samuel Eilenberg), pp. 63–76
- [11] Hurewicz Witold and H. Wallman, *Dimension Theory*, Princeton University Press, Princeton, New Jersey 1941
- [12] Ionicioiu Radu (1997), Building block for topology change in 3D, [gr-qc/9711069](#)
- [13] Johnson M., The combinatorics of n-categorical pasting, *Journal of Pure and Applied Algebra* **62** (1989) 211–225
- [14] Joyal André & Ross Street, Braided tensor categories, *Advances in Mathematics* **102** (1993) 20–78
- [15] Kerler Thomas (1998), Bridged links and tangle presentation of cobordism categories, [math.GT/980614](#)
- [16] Gordon R., A. J. Power, and Ross Street (1995), Coherence for Tricategories, *Memoirs of the American Mathematical Society*, Number 558, **117**
- [17] Leinster Tom (1998), Basic bicategories, [math.CT/9810017](#), [math.QA/0002180](#)
- [18] Oziewicz Zbigniew, Calculus of variations for multiple valued functionals, *Reports on Mathematical Physics* **31** (1) (1992) 85–90
- [19] Oziewicz Zbigniew and Guillermo Arnulfo Vázquez Coutiño, Categorical analysis, December 1999. Submitted to ‘Miscellanea Algebraica’, Waldemar Korczyński and Adam Obtułowicz, Editors, Akademia Świętokrzyska, Kielce, Poland.
- [20] Power A. J., A 2-categorical pasting theorem, *Journal of Algebra* **129** (1990) 439–445

- [21] Power A. J., An n-categorical pasting theorem, Springer Lectures Notes in Mathematics **1488**, New York
- [22] Power A. J., Why tricategories?, Info. Comp. **120** (1995) 251–262
- [23] Sawin Stephen, Links, quantum groups and TQFT's, Bulletin (New Series) AMS **33** (1996) 413
- [24] Simpson Carlos, A closed model structure for n-categories, internal Hom, n-stacks and generalized Seifert - Van Kampen, alg-geom/9704006
- [25] Simpson Carlos, Homotopy types of strict 3-grupoids, math.CT/981059
- [26] Steiner Richard, Pasting in multiple categories, Theory and Applications of Categories **4** (1) (1998) 1–36
- [27] Stong R. E., Notes on Cobordism Theory, Princeton University Press, 1968
- [28] Street Ross, The role of Michael Batanin's monoidal globular categories, Contemporary Mathematics **230** (1998) 99–116
- [29] Witten Edward, On quantum gauge theories in two dimensions, Communications in Mathematical Physics **141** (1991) 153
- [30] Wolff H., V-category and V-graph, Journal of Pure and Applied Algebra **4** (1974) 123–135
- [31] Yetter David N., Quantum groups and representations of monoidal categories, Mathematical Proceedings of Cambridge Philosophical Society **108** (1990) 261–290